

AD-A105 658

NORTH CAROLINA UNIV AT CHAPEL HILL DEPT OF STATISTICS
COMPACT COVARIANCE OPERATORS.(U)
AUG 81 C R BAKER, I W MCKEAGUE

F/6 12/1

N00014-75-C-0491

UNCLASSIFIED

NL

1 of 1
AD-A105 658



END
DATE
FILMED
11-81
DTIC

① LEVEL II

AD A105658

COMPACT COVARIANCE OPERATORS[†]

by

Charles R. Baker and Ian W. McKeague

ABSTRACT

Let B be a real separable Banach space and $R: B^* \rightarrow B$ a covariance operator. All representations of R in the form $\sum e_n \otimes e_n$, $\{e_n, n \geq 1\} \subset B$, are characterized. Necessary and sufficient conditions for R to be compact are obtained, including a generalization of Mercer's theorem. An application to characteristic functions is given.

KEY WORDS AND PHRASES: Covariance operators, compact operators, probability in Banach spaces.

AMS 1970 Subject Classifications: Primary 60B05, 47B05, 60B10.

DTIC
ELECTE
OCT 16 1981
S B D

This research was supported by the Office of Naval Research under Contract N00014-75-C-0491.

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

81 10 15

182850

DTIC FILE COPY

1. Introduction

The study of covariance operators is a major component in the theory of probability measures on Banach spaces [10], [2], [14]. The covariance operator of a strong second-order measure is always compact [2]; however, the covariance operator of a weak second-order measure need not be compact. In this paper we first characterize series representations of covariance operators, and then give a set of necessary and sufficient conditions for a covariance operator to be compact. The classical Mercer's theorem [14] can be obtained as an immediate corollary. These results are then applied to extend a result of Prohorov and Sazanov [6] on relative compactness of probability measures from Hilbert space to Banach space.



2. Definitions and Notation

B is a real separable Banach space with norm $||\cdot||$ and topological dual B^* . A linear operator $R: B^* \rightarrow B$ is a covariance operator if R is symmetric and non-negative: $\langle Ru, v \rangle = \langle u, Rv \rangle$ and $\langle Ru, u \rangle \geq 0$, for all u, v in B^* . A probability measure μ on the Borel σ -field of B is said to be weak second-order if $\int_B \langle x, u \rangle^2 d\mu(x) < \infty$, for all u in B^* ; μ is strong second-order if $\int_B ||x||^2 d\mu(x) < \infty$. Every weak second order measure μ has a mean element m in B and a covariance operator $R: B^* \rightarrow B$ [9], [10], defined by

$$\langle m, v \rangle = \int_B \langle x, v \rangle d\mu(x)$$

$$\langle Ru, v \rangle = \int_B \langle x - m, u \rangle \langle x - m, v \rangle d\mu(x),$$

for all u, v in B^* . Strong second-order measures have compact covariances; the strong second order property is not necessary in order that μ have compact covariance.

For a covariance operator $R: B^* \rightarrow B$ it is well known [8], [1], that there exists a separable Hilbert space $H \subset B$ such that the natural injection

$j: H \rightarrow B$ is continuous and $R = jj^*$. H is the RKHS of R and is the completion of range (R) with respect to the inner product $\langle \cdot, \cdot \rangle_H$ defined by $\langle Ru, Rv \rangle_H = \langle Ru, v \rangle$.

I_H will denote the identity on H . For u, v in B^* , z in B (resp. in H), $(u \otimes v)(z) = \langle v, z \rangle u$ (resp., $\langle v, z \rangle_H u$). If T is any map $r(T) \equiv \text{range}(T)$. τ_C is the linear topology on B^* determined by a neighborhood base at zero of the form $V_{C, \epsilon}(0) = \{f \in B^*: \sup_{x \in C} \langle f, x \rangle^2 < \epsilon\}$ for all $\epsilon > 0$ and all compact sets $C \subset B$ (τ_C is the topology of uniform convergence on compact sets). For a given covariance operator $R: B^* \rightarrow B$, q_R is the real-valued quadratic functional on B^* defined by $q_R f = \langle Rf, f \rangle$. The notation $R = \sum_n e_n \otimes e_n$ for $\{e_n, n \geq 1\} \subset B$ means that the sequence $(\sum_1^N e_n \otimes e_n)$ converges to R in the strong operator topology: $\sum_1^N \langle e_n, f \rangle e_n \rightarrow Rf$ in the norm topology of B , for all f in B^* . $I_H = \sum_n e_n \otimes e_n$ has a similar interpretation. If $\{u_n, n \geq 1\}$ is any orthonormal basis for H , then $R = \sum j u_n \otimes j u_n$, [9]. K_R will denote the unit ball in H .

If μ is a probability measure on the Borel σ -field of B , its characteristic functional $\hat{\mu}$ is defined as

$$\hat{\mu}(x) = \int_B e^{i\langle x, y \rangle} d\mu(y), \text{ for } x \text{ in } B^*.$$

3. Representation of Covariance Operators.

In this section, R is an arbitrary covariance operator.

Theorem 1. $R = \sum_n e_n \otimes e_n$ if and only if $e_n = j v_n$, $v_n \in H$ for $n \geq 1$, and

$$I_H = \sum_n v_n \otimes v_n.$$

For	<input checked="" type="checkbox"/>
MAI	<input type="checkbox"/>
ed	<input type="checkbox"/>
tion	

Distribution/	
Availability Codes	
Avail and/or	
Dist	Special
A	

Proof. It suffices to show that the stated conditions are necessary for $R = \sum_n e_n \otimes e_n$. Suppose $R = \sum_n e_n \otimes e_n$, and fix e_k . Let $P_k = e_k \otimes e_k$. To show $e_k \in \text{range}(j)$, let (as in [3]) $D: r(j^*) \rightarrow B$ be defined by $Dj^*f = P_k f$. Then $\|Dj^*f\|^2 = \|P_k f\|^2 = \|e_k\|^2 \langle e_k, f \rangle^2 \leq \|e_k\|^2 \sum_n \langle f, e_n \rangle^2 = \|e_k\|^2 \langle Rf, f \rangle = \|e_k\|^2 \|j^*f\|_H^2$. Thus D can be extended to a continuous linear map from $\overline{r(j^*)} = H$ into B . From its definition, $Dj^* = P_k$, so $P_k = jD^*$ and thus $e_k \in \text{range}(j)$.

To see that $I_H = \sum v_n \otimes v_n$, where $jv_n = e_n$, $n \geq 1$, define $Q_N = \sum_1^N v_n \otimes v_n$. $Q_N = Q_N^*$ and $Q_N \geq 0$, so $Q_N^{1/2}$ exists. $\|Q_N^{1/2} j^*f\|_H^2 = \sum_1^N \langle f, e_n \rangle^2 + \|j^*f\|_H^2$, so that $\|Q_N^{1/2}\| \leq 1$ and $\|Q_N^{1/2} x\|_H \rightarrow \|x\|_H$ for all x in $r(j^*)$. Thus,

$$\begin{aligned} \|\sum_1^N (v_n \otimes v_n) j^*f - j^*f\|_H^2 &= \|Q_N j^*f - j^*f\|_H^2 \\ &\leq \|Q_N^{1/2} j^*f\|_H^2 + \|j^*f\|_H^2, \end{aligned}$$

which converges to zero as $N \rightarrow \infty$ for any fixed f in B^* . Thus, $\sum v_n \otimes v_n = I_H$ on $r(j^*)$, and the result follows by $\overline{r(j^*)} = H$. \square

Remark. Suppose E is a locally convex topological vector space, $R: E' \rightarrow E$ is a covariance operator, and $R = jj^*$, where $j: H \rightarrow E$ is the injection and H is the RKHS of R . R will have such a representation, for example, if E is separable and quasi-complete [8]. In this case, it is easily shown that Theorem 1 holds without modification.

The representation $I_H = \sum v_n \otimes v_n$ does not require that $\{v_n, n \geq 1\}$ be a CONS in H ; however, sufficient conditions for $\{v_n, n \geq 1\}$ to be a CONS in H can be given.

Proposition 1. Suppose $I_H = \sum v_n \otimes v_n$; the following are equivalent:

- (1) $\|v_k\|_H = 1$
- (2) $v_k \notin \overline{\text{sp}\{v_n, n \neq k\}}$
- (3) $v_k \perp \overline{\text{sp}\{v_n, n \neq k\}}$.

If any of the above conditions hold for all $k \geq 1$, then $\{v_n, n \geq 1\}$ is a CONS for H .

4. Compact Covariance Operators.

Theorem 2. Suppose $R = \sum e_n \otimes e_n$, $\{e_n, n \geq 1\} \subset B$. Let $\{v_n, n \geq 1\} \subset H$ be such that $e_n = jv_n$, $n \geq 1$. The following are equivalent:

- (1) R is compact;
- (2) j is compact;
- (3) $j[K_R]$ is compact in B ;
- (4) the series $\sum v_n \otimes jv_n$ converges uniformly in H on bounded subsets of B^* ;
- (5) $(\sum_{n=1}^N e_n \otimes e_n)$ converges to R uniformly in B on bounded subsets of B^* ;
- (6) q_R is w^* -continuous on bounded subsets of B^* ;
- (7) q_R is τ_C -continuous.

Proof. (1) \Rightarrow (2). Suppose $f_\alpha \rightarrow f$ in the w^* topology of B^* , where $\|f_\alpha\| \leq k$ for all α . Then $\|j^*f_\alpha - j^*f\|_H^2 = \langle R(f_\alpha - f), (f_\alpha - f) \rangle \leq 2k \|R(f_\alpha - f)\|_B$; since R is compact, $j^*f_\alpha \rightarrow j^*f$ in H [4, p. 486] and thus j is compact.

(2) \Rightarrow (3). j compact implies $j[K_R]$ is relatively compact in B . Since K_R is weakly compact in H and j is weakly continuous, $j[K_R]$ is weakly compact in B , and thus closed.

(3) \Rightarrow (2) by definition.

(2) \Rightarrow (4). By Theorem 1, $\sum v_n \otimes v_n = I_H$. Set $Q_N = \sum_{n=1}^N v_n \otimes v_n$. If $A \subset B^*$ is bounded, then $j^*[A]$ is compact; by Dini's theorem $\|Q_N^*x\|_H \uparrow \|x\|_H$ uniformly on $j^*[A]$. Hence $\|(Q_N - I)j^*x\|_H^2 \leq \|j^*x\|_H^2 - \|Q_N^*j^*x\|_H^2 \rightarrow 0$ uniformly on A .

(4) \Rightarrow (5), since j is continuous.

(5) \Rightarrow (1), since R is the uniform limit of compact operators.

(2) \Leftrightarrow (6). Follows from the fact that j is compact if and only if $j^*f_\alpha \rightarrow 0$ in the norm topology of H for all bounded generalized sequences (f_α) in B^* which are w^* convergent to zero [4, p. 486], and $q_R(f_\alpha) = \|j^*f_\alpha\|_H^2$.

(1) \Rightarrow (7). Suppose R is compact. Writing $C = j[K_R]$, C is compact in B .
 $q_R(f) = \langle Rf, f \rangle = \|j*f\|_H^2 = \sup_{x \in K} \langle j*f, x \rangle_H^2 = \sup_{x \in C} \langle f, x \rangle^2$. Thus q_R is τ_C -continuous at zero. τ_C -continuity of q_R follows from $q_R(f_\alpha) = q_R(f_\alpha - f) - q_R(f) + 2\langle Rf, f_\alpha \rangle$.

(7) \Rightarrow (1). Suppose q_R is τ_C -continuous. Using (6), R is compact if q_R is w^* continuous at 0 on bounded subsets of B^* . But B is separable so that the w^* topology on bounded subsets of B^* is metrizable and it suffices to consider sequences. Suppose $f_n \xrightarrow{w^*} 0$ and $\|f_n\| \leq k$. Let L be an arbitrary compact subset of B . Since $\{f_n\}$ is bounded in B^* the f_n are equicontinuous and uniformly bounded as continuous functions on L . Thus, by the Arzela-Ascoli Theorem [4, p. 266] $\{f_n\}$ is relatively compact as a subset of $C^{\mathbb{R}}(L)$. Thus since $f_n \xrightarrow{w^*} 0$, f_n converges to 0 uniformly on L . Therefore $f_n \xrightarrow{\tau_C} 0$ and $q_R(f_n) \rightarrow 0$. This completes the proof of Theorem 2. \square

Remarks. (1) Suppose $r: [0,1] \times [0,1] \rightarrow \mathbb{R}$ is continuous, symmetric and positive definite. For fixed $t \in [0,1]$, let $\pi_t(x) = x_t$ for x in $C[0,1]$; $\|\pi_t\| = 1$. A compact covariance operator $R: C^*[0,1] \rightarrow C[0,1]$ is defined by $[R\mu](t) = \int_0^1 r(t,s) d\mu(s)$ for any μ in $C^*[0,1]$ (by Arzela-Ascoli Theorem). Thus for $s, t \in [0,1]$, $\langle R\pi_t, \pi_s \rangle = r(t,s)$. The integral operator in $L_2[0,1]$, corresponding to the kernel r , has continuous orthonormal eigenvectors $\{y_n, n \geq 1\}$ and associated non-zero eigenvalues $\{\lambda_n, n \geq 1\}$; it is well known that $\{\lambda_n^{1/2} y_n, n \geq 1\}$ is a CONS in the RKHS H of R . Thus, from Theorem 2, $\sum_{n=1}^N \lambda_n y_n(t) y_n(s)$ converges uniformly to $r(t,s)$ for all t, s in $[0,1]$. This is the classical Mercer's Theorem [7, pp. 245-246].

(2) The fact that the unit ball of H is compact in B when R is compact was proved by Kuelbs [5] under the assumption that R is the covariance of a strong second-order measure.

5. Characteristic Functionals

Let Λ denote a family of probability measures on B (separable Banach) and $\hat{\Lambda}$ the corresponding family of characteristic functionals.

Theorem 3. Let B be a separable Banach space. Then the following are equivalent:

- a) There exists a topology τ on B^* such that for each family Λ of probability measures on B , $\hat{\Lambda}$ is equicontinuous in this topology if and only if Λ is relatively compact in the topology of weak convergence
- b) B is finite dimensional.

Proof. As in the Hilbert space case (see [6, Lemma 2]) τ_c is the weakest topology on B^* such that relative compactness of $\Lambda \Rightarrow$ equicontinuity of $\hat{\Lambda}$. Suppose that (a) holds. Then $\tau_c \subset \tau$ and τ_c equicontinuity of $\hat{\Lambda}$ implies relative compactness of Λ . Now let $R: B^* \rightarrow B$ be any compact covariance operator. Let $\{e_n\}$ be a CONS in the RKHS of R . Define μ_k to be the zero mean Gaussian measure on B with covariance operator $\sum_{n=1}^k e_n \otimes e_n$. Then $\{\hat{\mu}_k\}$ is τ_c equicontinuous by Theorem 2 and $\{\mu_k\}$ is relatively compact. Therefore R is the covariance of a Gaussian probability measure on B and by [9, Theorem 11] B is finite dimensional. \square

Theorem 3 extends a result of Prohorov and Sazonov [6] who proved it for Hilbert spaces.

REFERENCES

- [1] Baxendale, P. (1976). Gaussian measures on function spaces. Amer. J. Math. 98, 891-952.
- [2] Chobanjan, S.A. and Tarieladze, V.I. (1973). On the compactness of covariance operators. (Russian). Soobsc. Acad. Nauk Gruzin, 70, 273-276.
- [3] Douglas, R.G. (1966). On majorization, factorization, and range inclusion of operators in Hilbert space. Proc. Amer. Math. Soc. 17, 413-415.
- [4] Dunford, N. and Schwartz, J.T. (1966). Linear Operators I. Interscience, New York.
- [5] Kuelbs, J. (1976). The law of the iterated logarithm and related strong convergence theorems for Banach space valued random variables. École d'Été de Probabilités de Saint-Flour 1975, Lecture Notes in Mathematics 539, Springer-Verlag, Berlin.
- [6] Prohorov, Yu. V. and Sazonov, V.V. (1961). Some results associated with Bochner's theorem. Theory Prob. Appl. 6, 82-87.
- [7] Riesz, F. and Nagy, B. Sz. (1956). Functional Analysis, Blackie and Son, London and Glasgow.
- [8] Schwartz, L. (1964). Sous-espaces Hilbertiens d'espaces vectoriels topologiques et noyaux associés. J. Analyse Math. 13, 115-256.
- [9] Vakhania, N.N. and Tarieladze, V.I. (1978). Covariance operators of probability measures in locally convex spaces. Theory Prob. Appl. 23, 1-21.
- [10] Weron, A. (1976). On weak second order and Gaussian random elements. Lecture Notes in Mathematics 526, 263-272, Springer-Verlag, Berlin.

Department of Statistics
University of North Carolina
Chapel Hill, NC 27514

Department of Statistics
Florida State University
Tallahassee, FL 32306

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
AD A205658		
4. TITLE (and Subtitle) Compact Covariance Operators		5. TYPE OF REPORT & PERIOD COVERED TECHNICAL
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Charles R. Baker and Ian W. McKeague		8. CONTRACT OR GRANT NUMBER(s) ONR Contract N00014-75-C-0491
9. PERFORMING ORGANIZATION NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Statistics and Probability Program (Code 436) Arlington, VA 22217		12. REPORT DATE August 1981
		13. NUMBER OF PAGES 8
14. MONITORING AGENCY NAME & ADDRESS, if different from Controlling Office		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for Public Release -- Distribution Unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Covariance operators, compact operators, probability in Banach spaces.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Let B be a real separable Banach space and $R: B^* \rightarrow B$ a covariance operator. All representations of R in the form $\sum e_n \otimes e_n$, $\{e_n, n \geq 1\} \subset B$, are characterized. Necessary and sufficient conditions for R to be compact are obtained, including a generalization of Mercer's theorem. An application to		

DD FORM 1473

JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

20. characteristic functions is given.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

**DATA
FILM**